Math 371 (Differential Equations), Section 3 (Dmytro Savchuk) Fall 2009 Homework assignment 8

Note: I will not collect this assignment – just do it for your benefit. This is, perhaps, the most important HW assignment in the course. This is a preparational homework for the final that covers the topics that will be presented on the final. Here is a list of these topics:

- 1. First order equations using exactness or integrating factors
- 2. Euler-Cauchy problem
- 3. Orthogonal family of curves
- 4. 2×2 systems of linear equations with distinct roots for the polynomial
- 5. Power series with solution to 5th order
- 6. Second solution using reduction of order
- 7. Laplace

Solve the following problems

1. Check that the following differential equation is exact and solve it

$$(1 + e^{x}y + xe^{x}y)dx + (xe^{x} + 2)dy = 0$$

2. Make the following equation exact by multiplying by appropriate integrating factor and solve it

$$(2x^2 + y)dx + (x^2y - x)dy = 0$$

3. Solve the Euler-Cauchy equation

$$x^2y'' - 5xy' + 8y = 0$$

4. Solve the Euler-Cauchy equation

$$x^2y'' - 5xy' + 9y = 0$$

5. Solve the Euler-Cauchy equation

$$x^2y'' - 5xy' + 10y = 0$$

- 6. Find a family of curves orthogonal to the family $y = kx^4$.
- 7. Find a family of curves orthogonal to the family $y^2 = kx$.
- 8. Solve the system of differential equations

$$\begin{cases} y_1' = y_1 + 2y_2 \\ y_2' = y_1 + 2y_2 \end{cases}$$

9. Find the solution of differential equation 2y'' + xy' + y = 0 using power series. Approximate the solution by taking explicitly the first 6 terms in your power series (up to x^5).

- 10. Find the solution of differential equation $x^2y'' + 3y' xy = 0$ using power series. Approximate the solution by taking explicitly the first 6 terms in your power series (up to x^5).
- 11. Given that $f(x) = e^x$ is the solution to the differential equation xy'' (x+1)y' + y = 0 find the second linearly independent solution.
- 12. Find the Laplace transforms of the functions

(a)
$$f(t) = e^{-2t} \sin 2t + e^{3t}t^2$$

(b) $f(t) = sin^2(4t)$
(c) $f(t) = t^2u(t-4)$
(d) $f(t) = \begin{cases} t^2, & t < 2, \\ \sin(t), & 2 < t < 4, \\ e^{-t}, & t > 4. \end{cases}$

13. Find the inverse Laplace transforms of the functions

(a)
$$F(s) = e^{-s} \frac{s}{s^2 + 4}$$

(b) $F(s) = \frac{s + 1}{s^2 + 2s + 10}$
(c) $F(s) = \frac{5s^2 + 34s + 53}{(s + 3)^2(s + 1)}$
(d) $F(s) = \ln \frac{s + 2}{s - 5}$

14. Use Laplace transform to solve the initial value problem

$$y' - y = u(t - 1),$$
 $y(0) = 0$

15. Use the Laplace transform to solve the initial value problem

$$y' - y = u(t - 1),$$
 $y(0) = 0$

16. Use the Laplace transform to solve the initial value problem

$$y' - 2y = u(t - 2), \qquad y(0) = 1$$

17. Use the Laplace transform to solve the initial value problem

$$y'' + 9y = 3\delta(t - \pi), \qquad y(0) = 1, \quad y'(0) = 0$$

18. Use the Laplace transform, the method of undetermined coefficients and the variation of parameters to solve initial value problem

$$y'' - 2y' + 5y = -8e^{-t}, \qquad y(0) = 2, \quad y'(0) = 12$$